Infant CPR Skills Testing Checklist



Student Name	Date of Test	
Scenario: "While you are pushing a baby in a stroller at the park, you notice something is wrong with the baby. You do not have a phone nearby. You ensure that the scene is safe and take the baby out of the stroller. Demonstrate what you would do next."		
Assessment and Activation ☐ Checks responsiveness ☐ Shouts for help ☐ Checks	breathing	
Once student shouts for help, instructor says, "No one is around to help." Cycle 1 of CPR (30:2) *CPR feedback devices preferred for accuracy		
Infant Compressions ☐ Performs high-quality compressions*: • Placement of 2 fingers in the center of the chest, just below the nipple line • 30 compressions in no less than 15 and no more than 18 seconds • Compresses at least one third the depth of the chest, about 1½ inches (4 cm) • Complete recoil after each compression Cycle 2 of CPR (repeats steps in Cycle 1) Only chest Gives 30 high-quality compressions ☐ Gives 2 effective Gives 30 high-quality compressions ☐ Gives 30 high-quality c	breaths eck box if step is successfully perfo	econd breath 10 seconds
Instructor says, "You have just completed 5 sets of 30 compressions and 2 breaths."		
Activates Emergency Response System (9-1-1) Uerbalizes the need to leave to phone 9-1-1		
STOP TEST		
 Instructor Notes Place a ✓ in the box next to each step the student completes so If the student does not complete all steps successfully (as indicate remediation. Make a note here of which skills require remediation. 	ed by at least 1 blank check box), the s	l l
Test Results Check PASS or NR to indicate pass or needs re	emediation:	PASS NR
Instructor Initials Instructor Number	Date	

Infant CPR Skills Testing Critical Skills Descriptors

- 1. Assesses infant and activates emergency response system (this *must* precede starting compressions) within a maximum of 30 seconds. After determining that the scene is safe:
 - Checks for responsiveness by tapping and shouting
 - · Shouts for help
 - Checks for no breathing or no normal breathing (only gasping)
 - Scans from the head to the chest for a minimum of 5 seconds and no more than 10 seconds
- 2. Cycle 1: Performs high-quality chest compressions (initiates compressions immediately after recognition of cardiac arrest)
 - · Correct placement of fingers in center of chest
 - 2 fingers just below the nipple line
 - Compression rate of 100 to 120/min
 - Delivers 30 compressions in 15 to 18 seconds
 - Compression depth and recoil—compress at least one third the depth of the chest, about 11/2 inches (4 cm)
 - Use of a commercial feedback device/manikin is highly recommended
 - Complete chest recoil after each compression
- 3. Cycle 1: Provides 2 breaths by using a barrier device
 - · Opens airway adequately
 - Uses a head tilt-chin lift maneuver
 - · Delivers each breath over 1 second
 - · Delivers breaths that produce visible chest rise
 - · Avoids excessive ventilation
 - Resumes chest compressions in less than 10 seconds
- 4. Cycle 2: Performs same steps for compressions and breaths as in Cycle 1
- 5. Cycle 3: Performs same steps for compressions and breaths as in Cycle 1